

# Wildberry Farms Vaccination Guidelines

All cattle will be required at weaning to have vaccinations (primary and booster), 2-4 weeks apart. A Modified Live Vaccine is required for the booster. All vaccinations should be given in the neck according to BQA guidelines. Male calves must be castrated and healed, females must be open and all calves must be polled or dehorned.

Producer \_\_\_\_\_ #Head \_\_\_\_\_

## Primary Vaccinations (3-4 weeks pre-weaning recommended)

BVD, IBR, PI-3 BRSV

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## 7-Way Clostridial (blackleg) + Haemophilis Somnus

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## Mannheimia (Pasteurella) Haemolytica bacterin/leukotoxioid

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## Implants: Please No Implants at Weaning.

Cattle implanted as young calves are acceptable, however, we cannot accept calves implanted at weaning.

## Dewormed (internal parasites, lice & grubs)

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## Booster Vaccinations (2-4 weeks following primary vaccinations)

BVD, IBR, PI-3 BRSV (Modified Live Viral Vaccine Required)

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## 7-Way Clostridial (blackleg) + Haemophilis Somnus

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## Mannheimia (Pasteurella) Haemolytica bacterin/leukotoxioid

Product Name & Brand \_\_\_\_\_

Serial # \_\_\_\_\_ Lot# \_\_\_\_\_

Date Administered \_\_\_\_\_ Expiration Date \_\_\_\_\_

## Recommendations

Bovi-Shields Gold 5; Pyramid 4  
Pyramid 5; Vista 5 SQ

Vision 7 Somnus; Vision 8 Somnus  
Ultrabac 7/Somubac;  
Bar-Vac 7/Somnus; One Shot Ultra 7

One Shot; Presponse SQ;  
Presponse HM; Pulmo-Guard PHM-1;  
Pulmo-Guard PHM; Once PMH  
(Copy the information if the vaccine is included with another vaccine dose)

No Implants at Weaning

Dectomax; Cydectin;  
Ivomec

Booster same as above

Booster same as above

Booster if primary was not a single dose product

All cattle should be weaned a minimum of 30 days with boosters given at least 14 days prior to delivery. A veterinarian is not required to participate in the program but we do recommend consulting your veterinarian before starting a modified live viral pre-weaning vaccination protocol.

I believe the above information is true and accurate.

Producer Signature \_\_\_\_\_ Date \_\_\_\_\_